# Re-blessing: Study of ttbar Production Mechanisms

Analysis Overview

Changes to the previously blessed results

Plots and Results to Bless

Shabnaz Pashapour, Pekka K. Sinervo University of Toronto

Top Group Meeting July 19, 2007

## Analysis Overview M

- The goal is to measure  $\sigma_{(gg o t \bar{t}\,)}/\sigma_{(p\bar{p} o t \bar{t}\,)}$ 
  - √ Test of pQCD
  - √ High x gluon distribution
  - ✓ Unknown sources of physics beyond the SM
- Low pt track multiplicity
- Data-driven method
- Dijet and W+n jet samples as calibration

#### Track multiplicity


#### defTracks

$$✓$$
 p<sub>T</sub> 0.3 – 2.9 GeV/c<sup>2</sup>

$$\sqrt{|\eta|} \le 1.1$$


- Matched to the event vertex
  - √ 3cm
- Away from jets
  - ✓  $\Delta$ R=0.6, corE<sub>T</sub>≥15 GeV
  - ✓  $\Delta$ R=0.4, 6 ≤ corE<sub>T</sub>< 15 GeV
- Correct for area differences
- Correct for remaining contribution of high E<sub>T</sub> jets
  - $\checkmark$  0d: 0.90  $\pm$  0.03
  - $\checkmark$  0h: 0.97  $\pm$  0.04
  - $\checkmark$  0i: 0.96 ± 0.04

#### Track if no magnetic field exists



Track in magnetic field

#### Jet of 0.4 and its annuli



# Analysis Overview II

- Correlation between <N<sub>trk</sub>> and <N<sub>g</sub>>
  - ✓ MC calculations to find <N<sub>g</sub>> in a sample
- W+0 jet sample, almost purely qq
- dijet sample with leading jet Et of 80-100 GeV, large gluon content
- Binned likelihood fit with two free parameters

$$N[f_gF_g^{norm}+(1-f_g)F_q^{norm}]$$

# Changes to the Previously Blessed Results

#### Changes in W+n jet/ttbar event selection


- Bugs that are fixed
  - ✓ TCEM, TCMUP and TCMX triggers had been assumed to be fired.
    - 8 TCMUP ttbar candidates
  - $\checkmark$  For the tight jets, the event  $\eta$  was used instead of the detector  $\eta$ .
    - 3 ttbar candidates
- QCD rejection cut
  - ✓ We had required  $\Delta φ$  of the missing Et and the leading tight jet to be between 0.5-2.5 rad, if missing Et < 30 GeV
    - To be consistent with the background estimates, we removed it.
      16 ttbar candidates
- Now observe 240 tagged lepton+jet ttbar candidate events, instead of 229.

#### QCD background in W+0 jet

 Previously estimated 1% had assumed W mass constraint, we do not and so now estimate it to be


$$N_D^{\text{bkg}}/N_D = (N_A/N_B)(N_C/N_D) = (4.9\pm0.4)\%$$


- Electrons and muons are different separately, use for systematic uncertainties
- Gluon-content change by less than 0.5%



#### Effects and implications...

- Negligible change in parameterizations.
- Bugs and W+0 jet QCD background has negligible effects.
- Removing QCD veto is important as previously we were doublecounting background.
- Use previous systematics
  - ✓ checked W+0 gluon composition, no change.








# Correlation between <Ng> and <Ntrk>


Sample	MC <n<sub>g&gt;</n<sub>	Data <n<sub>trk&gt;</n<sub>
W+0 jet	0.05 ±0.10	10.22±0.01
W+1 jet	0.97 ±0.10	11.25 ±0.03
W+2 jets	1.38 ±0.10	11.47 ±0.06
80-100 GeV	1.72 ±0.10	12.33 ±0.02
100-120 GeV	1.62 ±0.10	12.10 ±0.02
120-140 GeV	1.44 ±0.10	11.96 ±0.04



#### using the fit to find < Ng > for other samples < Ntrk >

Sample	MC prediction	Fit result
140-160 GeV	1.26 ±0.04	1.39 <sup>+0.06</sup> <sub>-0.05</sub>
160-180 GeV	1.13 ±0.04	1.23±0.05
180-200 GeV	0.99 ±0.07	1.08+0.05
200-220 GeV	0.92 ±0.10	0.88+0.04
220+ GeV	0.67 ±0.10	0.65+0.05

#### Parameterization



#### Fit and MC values for different calibration samples

Sample	$f_{\!\scriptscriptstyle g}$ from the fit	MC prediction
80-100 GeV	$0.734 \pm 0.004$	0.73 ±0.02
100-120 GeV	$0.688 \pm 0.005$	0.69 ±0.02
120-140 GeV	0.659 ± 0.010	0.63 ±0.03
140-160 GeV	$0.627 \pm 0.005$	0.57 ±0.03
160-180 GeV	$0.573 \pm 0.005$	0.52 ±0.03
180+ GeV	$0.492 \pm 0.005$	0.42 ±0.05

#### Estimating gluon-rich fraction in background

Sample	$f_{g}$ no tag	$f_g$ _tagged
W+1 jet	0.41 ± 0.01	$0.56 \pm 0.05$
W+2 jet	0.51 ± 0.02	$0.42 \pm 0.08$
W+3 jet	0.56 ± 0.04	0.44 ± 0.12
Extrapolated W+4+ jet, $(f_g^{LF})$ $(f_g^{HF})$	$0.72 \pm 0.05$	0.27 ± 0.19
LF fraction in background $(f_b^{LF})$	-	0.55 ± 0.11
HF fraction in background $(f_b^{HF})$	-	$0.45 \pm 0.09$

• We calculate  $f_g^{bkg}$  assuming Gaussian distributions for the variables used in the following equation using the above values

$$f_g^{bkg} = f_b^{LF} f_g^{LF} + f_b^{HF} f_g^{HF}$$

- We find  $f_g^{bkg}$  = 0.53  $\pm$  0.09 (modeling)  $\pm$  0.09 (nonW HF/LF composition)
- HF background is anything that can have a real tag (Wc, Wcc, Wbb, Single Top and half of nonW) and the rest is what we consider LF

## gg and aq to ttbar Acceptance

	gg→tt, ≥4 jets	qq→tt, ≥4 jetsUpdated
Updated A <sub>tagged</sub>	0.141 ±0.005	0.115 ±0.004
Previous A <sub>tagged</sub>	0.099 ±0.003	0.088 ±0.003

Used ttop75 PYTHIA MC Sample

## Systematic uncertainties-l

Uncertainties affecting track multiplicity

✓ Change the central values and observe the

changes in relevant variables

changes in relevant variables	$f_{g}$	$f_g^{\ bkg}$
Track/jet correction	±0.051	±0.001
Low jet E <sub>⊤</sub> cut	±0.021	±0.035
Dijet qq→qq fraction	±0.002	±0.019
W+0 jet $f_g$	±0.039	±0.007
nonW LF/HF composition	-	±0.057
Modeling the $f_g^{\ bkg}$ distribution	-	±0.089
Total	±0.07	±0.11

# Systematic uncertainties-11

• Uncertainties due to  $f_g$ ,  $f_g^{bkg}$  and  $f_b$ 

	$f_{g}^{tt}$
$f_{g}$	±0.08
$f_{g}^{bkg}$	±0.02
$f_{b}$	±0.01
Total	±0.08

## Systematic uncertainties-111

Uncertainties due to f<sub>g</sub><sup>tt</sup> and acceptances

	$\sigma(gg \rightarrow tt)/\sigma(pp \rightarrow tt)$
$f_{g}^{\;tt}$	±0.067
$A_{gg o tt}/A_{qq o tt}$	±0.004
Total	±0.07

#### Result

Using the fit result

$$f_g^{W+\geq 4} = 0.15 \pm 0.14 (stat) \pm 0.07 (syst)$$

• and the values we found, and a background fraction of  $(13 \pm 2)\%$ , we get

$$f_g^{tt} = 0.09 \pm 0.16 (stat) \pm 0.08 (syst),$$

• and using  $A_{gg \to tt}/A_{qq \to tt}$  = 1.23 ± 0.06, we find

$$\frac{\sigma(gg \to t\bar{t})}{\sigma(p\bar{p} \to t\bar{t})} = 0.07 \pm 0.14(stat) \pm 0.07(syst)$$

